

## AfriDX

COVID-19 diagnostics for Africa

# Silica Purification of Engineered Boon2-BST

Cambridge Analytical Biotechnology Group  
Department of Chemical Engineering and Biotechnology  
University of Cambridge



This project was funded by EDCTP and the European Union's Horizon 2020 research and innovation program under grant number RIA2020EF-2918

### Preliminary note to user:

- The following protocol was designed for purifying Boon2-BST from BL21 (DE3) cells
- For greater lysis efficiency, probe sonication can be used.

### Materials:

- An induced *E. coli* cell pellet
- Lysozyme
- 1X PBS (pH 7.2)
- Microcentrifuge Tubes
- Pipettes
- 60 $\mu$ m silica gel

### Protein Purification Protocol for Engineered Boon2-BST:

**Step 1:** Suspend the cell pellet with 1 mg/mL of lysozyme in 1X PBS (pH 7.2). The final volume of the suspension is dependent on the volume of the bacteria culture used to form the pellet. For this protocol, a ratio of 1 mL of lysozyme per 10 mL *E. coli* culture is used.

**Step 2:** Incubate the lysis solution at room temperature for 30 minutes.

**Step 3:** Prepare a solution of 60 $\mu$ m silica gel with a concentration of 25 mg/mL using 1X PBS (pH 7.2). It should be noted that vortexing and inverting the tube/container can assist in suspending the silica gel. In addition, this solution can be prepared and set aside during the lysis incubation step.

**Step 4:** After the lysis incubation, aliquot the cell lysis samples into microcentrifuge tubes.

**Step 5:** Centrifuge the lysis samples for 30 minutes at 13 krpm.

**Step 6:** Collect the cell lysate supernatant from the centrifuged samples. The solution should be pink in colour.

**Step 7:** Prepare the silica for immobilisation by first preparing microcentrifuge tubes with 20 mg of 60 $\mu$ m silica gel (0.8 mL from the stock prepared in **Step 3**).

**Step 8:** Centrifuge the prepared silica at 7.5 krpm for 5 minutes.

**Step 9:** After centrifugation, remove the supernatant, leaving only silica gel in the tube.

**Step 10:** Add the cell lysate directly to the silica, at a ratio of 1 mL of cell lysate to 20 mg of silica gel.



**Step 11:** Suspend the silica in the cell lysate using a vortexer, and afterwards lay the tube flat on its side to spread out the settling microparticles.

**Step 12:** Incubate the silica with the cell lysate for 30 minutes at room temperature. To ensure even absorption across the silica microparticles, vortex the sample every 10 minutes.

**Step 13:** After silica immobilisation, first vortex the sample and then centrifuge at 7.5 krpm for 5 minutes.

**Step 14:** Remove the supernatant, leaving behind silica with immobilised Boon2-BST. The silica itself should be visibly pink, while the supernatant is colourless.

**Step 15:** Perform 2 rounds of washing on the silica immobilised Boon2-BST.

**Washing Steps:**

- Add 1 mL of 1X PBS (pH 7.2)
- Suspend the immobilised protein with a vortex
- Centrifuge the protein at 7.5 krpm for 5 minutes
- Removes the “wash” buffer

**Step 16:** Suspend the immobilised Boon2-BST in 1X PBS (pH 7.2) to its working concentration. A ratio of 5µL of buffer to 1 mg of silica gel is used.

**Step 17:** Collect and combine the immobilised Boon2-BST. The working stock can be stored at 4°C both short and long term.